Running Head: Spatial Frequency Tuning in Object Representations

IN PRESS: JOURNAL OF COGNITIVE NEUROSCIENCE

Spatial frequency tuning reveals interactions between the dorsal and ventral visual systems

Bradford Z. Mahon^{1,2,3}, Nicholas Kumar¹, and Jorge Almeida^{4,5,6}

¹Department of Brain and Cognitive Sciences, University of Rochester, Rochester, NY
 ²Department of Neurosurgery, University of Rochester, Rochester, NY
 ³Center for Visual Science, University of Rochester, Rochester, NY, USA
 ⁴Faculty of Psychology and Education Sciences, University of Coimbra, Portugal
 ⁵School of Psychology, University of Minho, Portugal
 ⁶ Faculty of Psychology, University of Lisbon, Portugal

Document Size: 27 Pages (~6,548 words) 6 Figures Abstract: 247 words Introduction: 726 words Discussion: 1354 words

Address for Correspondence Bradford Z. Mahon Meliora Hall University of Rochester Rochester, NY. 14627-0268 mahon@rcbi.rochester.edu

Abstract

It is widely argued that the ability to recognize and identify manipulable objects depends on the retrieval and simulation of action-based information associated with using those objects. Evidence for that view comes from functional Magnetic Resonance Imaging (fMRI) studies that have reported differential Blood Oxygen Level Dependent (BOLD) contrast in dorsal visual stream regions when participants view manipulable objects compared to a range of baseline categories. An alternative interpretation is that processes internal to the ventral visual pathway are sufficient to support the identification of manipulable objects, and that the retrieval of object-associated use information is contingent on analysis of the visual input by the ventral stream. Here we sought to distinguish these two perspectives by exploiting the fact the dorsal stream is largely driven by magnocellular input, which is biased toward low spatial frequency (LSF) visual information. Thus, any tool-selective responses in parietal cortex that are driven by high spatial frequencies (HSF) would be indicative of inputs from the ventral visual pathway. Participants viewed images of tools and animals containing only low, or only high, spatial frequencies during fMRI. We find an internal parcellation of left parietal 'tool-preferring' voxels: inferior aspects of left parietal cortex are driven by HSF information and have privileged connectivity with ventral stream regions that show similar category preferences, while superior regions are driven by LSF information. Our findings suggest that the automatic activation of complex object-associated manipulation knowledge is contingent on analysis of the visual input by the ventral visual pathway.

Spatial Frequency Tuning in Object Representations

A basic distinction within primate vision is between a dorsal visual objectprocessing stream that projects from primary visual cortex to posterior parietal cortex and a ventral visual object-processing stream that projects from primary visual cortex to ventral temporal cortex (Ungerleider and Mishkin, 1982; Felleman and Van Essen, 1991; Goodale, 1992). The classic understanding of the dorsal visual system is that it computes spatial and volumetric properties from the visual input to support grasping, locomotion and eye movements, while ventral pathways extract object identity across variation in orientation, distance, and size. Lesions to dorsal stream regions can lead to visuomotor impairments leaving object identification unaffected, whereas ventral stream lesions can impair object identification while sparing visuomotor abilities (Goodale and Milner, 1992; Pisella, Binkofski, Lasek, Toni, and Rossetti, 2006).

One difference between the types of visual information that are processed within the ventral and dorsal streams originates in two types of retinal ganglion cells. Midget ganglion cells are sensitive to mid-to-high spatial frequencies (HSF) and project information through parvocellular nerve pathways, whereas parasol ganglion cells are sensitive to low spatial frequencies (LSF) and relay information through magnocellular nerve pathways (Derrington et al., 1984; Livingstone and Hubel, 1988; Tootell et al., 1988). While parvocellular and magnocellular information mixes already in V1 (Livingstone and Hubel, 1988; Maunsell et al., 1990; Merigan and Maunsell, 1993), there is an asymmetry in the way the two channels of information are passed on to the ventral and dorsal streams (Merigan and Maunsell, 1993): the ventral visual pathway receives both magnocellular and parvocellular inputs (Ferrera et al., 1992), while the

processing of manipulable objects). If the assumption is accepted that concept retrieval is a necessary step in object identification (the denial of which would seem untenable) then it follows that, on such accounts, object identification entails motor retrieval.

The logic of the current investigation is to use spatial frequencies to gain leverage on the critical issue of whether the retrieval of object-associated manipulation knowledge is contingent on analysis of the visual input by the ventral object processing stream. As noted above, there is a strong bias against high spatial frequency information being processed by the dorsal object processing stream. Thus, in the measure to which tool preferences are observed in parietal cortex for stimuli containing only high spatial frequencies, then it follows that at least some dorsal stream object representations are contingent on analysis of the visual input by ventral visual structures. If that were the case, then the core assumption of the Embodied Cognition Framework as it applies to manipulable object identification would be undermined, and a theory of object recognition would have to explore how activation of some motorrelevant information about objects is contingent on ventral stream processes.

Materials and Methods

Participants. Forty participants were tested (mean age = 20.1 ± 1.2 yrs standard deviation). Twenty-five participants completed Experiment 1 and 15 Experiment 2. All participants had normal or corrected to normal eyesight and no history of neurological disorders. All guidelines and requirements of the University of Rochester's Research Subjects Review Board were followed for participant recruitment and experimental procedures.

Experimental Stimuli. Stimuli were black-and-white pictures of animals and tools on a grey background. Thirty-two animal and 32 tool objects were used, with two exemplars for each object (total of 128 pictures). The words corresponding to the tool and animal names were matched on length in letters (means, standard deviations in parenthesis: Animals = 5.5 (2.0), Tools = 5.9 (2.4); t < 1), lexical frequency using the CELEX database (Animals = 22.8 (30.6), Tools = 20.2 (37.2), t < 1), and concept familiarity using the MRC Online Psycholinguistic Database (Animals = 519.8 (33.9), Tool = 519.3 (62.4), t < 1). The pictures were enclosed in a 200 by 200 square pixel frame, and subtended $\sim 4^{\circ}$ of visual angle (viewing angle ~ 47 pixels per degree). The spatial frequency content of the original broadband images was filtered to create high and low spatial frequency filtered versions of the pictures, using Matlab (Matlab 2009). Briefly, images were Fourier transformed into frequency space, the target frequencies were filtered, and the filtered images transformed back to their original space through an inverse Fourier-transform. To create high-spatial frequency items, we used a high-pass cut-off that was higher than 7 cycles per degree, and to create the low-spatial frequency items we filtered the pictures using a low-pass cut-off that was lower than 1 cycle per degree. Average luminance of the pictures did not differ between the three spatial frequency conditions (Broad-band, HSF, LSF; mean luminance across all stimuli was 132.62, 132.60 and 132.62 on a 255 gray-level scale for Broad-band, HSF, and LSF images, respectively; F (2,381) < 1). Moreover, average luminance within each spatial frequency condition did not differ between the two semantic categories (Broad-band: 131.60 and 133.64 for animal and tools, respectively, t(63) = 1.515, p = .135; HSF images: 131.59 and 133.63 for animal and tools, respectively, t(63) = 1.524, p = .133;

LSF images: 131.60 and 133.65 for animal and tools, respectively, t(63) = 1.515, p = .135).

General Procedure. 'A Simple Framework' (Schwarzbach, 2011) was used to control stimulus presentation and response collection in Psych toolbox in MATLAB on a MacPro running windows (Brainard, 1997; Pelli, 1997). Stimuli were back projected (temporal resolution = 120Hz) on a screen that participants viewed with a mirror attached to the head coil.

Experiment 1. Participants viewed the tool and animal stimuli passively (no response) in a mini-block design. Within each 16 second miniblock, 32 animals or 32 tools were presented (Duration = 500ms, ISI = 0). Miniblocks of stimuli were separated by 16 seconds of fixation. Each exemplar (n = 2) was presented twice per run resulting in 4 miniblocks of animals and 4 of tools for each run. Baseline conditions consisted of scrambled versions of the same stimuli, creating 'scrambled animal' and 'scrambled tool' conditions that preserved (but randomly displaced) local low level visual information (scrambling kernel = 32 pixels). All participants completed two runs of the protocol in Experiment 1 (each run lasted ~9 minutes).

Experiment 2. Experiment 2 followed the same design and procedure as Experiment 1 except for the inclusion of two additional conditions: HSF and LSF images of tools and animals, both intact and scrambled. As in Experiment 1, each run contained a balanced experimental design (each run lasted ~7.5 minutes). Participants completed between 5 and 8 runs.

MRI Parameters. Whole brain BOLD imaging was conducted on a 3-Tesla Siemens MAGNETOM Trio scanner with a 32-channel head coil at the Rochester Center for Brain Imaging. High-resolution structural T1 contrast images were acquired using a magnetization prepared rapid gradient echo (MPRAGE) pulse sequence at the start of each session (TR = 2530 ms, TE = 3.44 ms flip angle = 7 degrees, FOV = 256 mm, matrix = 256 x 256, 1x1x1mm sagittal left-to-right slices). An echo-planar imaging pulse sequence was used for T2* contrast (TR = 2000 ms, TE = 30 ms, flip angle = 90 degrees, FOV = 256 mm, matrix 64 x 64, 30 sagittal left-to-right slices, voxel size = 4x4x4mm). The first 6 volumes of each run were discarded to allow for signal equilibration.

fMRI Data analysis. fMRI data were analyzed with the *Brain Voyager* software package (Version 2.1) and in-house scripts drawing on the BVQX toolbox for MATLAB. Preprocessing of the functional data included, in the following order, slice time correction (sinc interpolation), motion correction with respect to the first volume of the first functional run, and linear trend removal in the temporal domain (cutoff: 2 cycles within the run). Functional data were registered (after contrast inversion of the first volume) to high-resolution de-skulled anatomy on a participant-by-participant basis in native space. For each participant, echo-planar and anatomical volumes were transformed into standardized (Talairach and Tournoux, 1988) space. Functional data were smoothed at 6mm (1.5 voxels) at FWHM, and interpolated to 3x3x3mm voxels.

The general linear model was used to fit beta estimates to the events of interest. The first derivative of 3D motion correction from each run were added to all models as regressors of no interest to attract variance attributable to head movement. All analyses treated subjects as a random factor, and there were thus 24 and 14 degrees of freedom in the group-level analyses in Experiments 1 and 2, respectively. Experimental events

Spatial Frequency Tuning in Object Representations

were convolved with a standard 2 gamma hemodynamic response function. In Experiment 1 there were 6 regressors: the two by two design of first/second presentation of images (factor not analyzed herein) and the category of the stimulus (tools and animals), as well as the scrambled versions of tools and animals. In Experiment 2, there were 12 predictors of interest, corresponding to the orthogonal crossing of the following three factors: Spatial Frequency (HSF, LSF, Broad-band), Category (Tool, Animal), and Image type (Intact, Scrambled).

All functional connectivity analyses were time course based and used the time series from the entire run. Time courses were extracted from preprocessed functional data, that had also been regressed with the outputs from motion correction (change in head position across volumes) and the global mean time-course from the whole brain. All connectivity was then computed over the residuals from that model. Whole brain functional connectivity maps were computed with a mask fit to the deskulled talairached anatomy. Whole brain maps were computed on a run-by-run basis, and the run-specific r maps were then averaged (within voxel) for each subject. Group level statistics were computed using a one-sample ttest against zero within each voxel across subjects.

Results

Inferior-to-Superior Organization by Spatial Frequency Within Parietal Cortex

In Experiment 1 we localized tool-preferring regions in the dorsal and ventral streams. Participants viewed broad-band images of tools and animals arranged in category-homogenous miniblocks of 16 seconds (32 stimuli, each presented for 500ms; See Figure 1 for examples of the stimuli). Replicating previous studies (e.g., Chao and Martin, 2000; Noppeney et al., 2006; Mahon et al., 2007), viewing images of tools

(compared to viewing animals) elicited differential activation in three left hemisphere regions—left parietal cortex, the left middle temporal gyrus, and the left fusiform gyrus (FDR, q < .05). Figure 2A shows a detail of the left parietal activation for tools compared to animals. This result demonstrates that the current set of materials generates the same topography of tool-preferring voxels as previous work.

In Experiment 2 we tested whether the left parietal tool-preferring region could be parcellated according to whether the tool-preferences were carried by the low or the high spatial frequency information in the images. Tool preferences in parietal cortex were determined separately for HSF and LSF images, by contrasting HSF tools against HSF animals, and separately, LSF tools against LSF animals. The results (Figure 2B) indicate that tool preferences for HSF images are biased toward the inferior aspect of parietal tool-preferring cortex, while tool preferences for LSF images are biased toward the superior aspect of left parietal tool preferring cortex. Importantly, the contrast maps showing tool-preferences in parietal cortex for HSF and LSF and LSF stimuli are independent; thus, there is no reason, in principle, why the two category-contrasts could not lead to entirely overlapping populations of voxels (see also Figure 3 for overlap of the contrast maps from Experiments 1 and 2).

The finding that there is an inferior-to-superior organization within left parietal tool-preferring voxels according to the spatial frequency content of the images can be directly tied to what is known about the behavioral impairments that attend selective lesions to inferior or superior parietal regions. Optic ataxia, an impairment for reaching and/or grasping objects is classically associated with lesions to posterior and superior parietal structures (Goodale, 1992; Pisella et al., 2006). Importantly, optic ataxia is not

necessarily associated with difficulties for manipulating objects according to their function (once patients are able to prehend the objects). In contrast, an impairment for manipulating objects correctly according to their function, termed apraxia of object use, is classically associated with lesions to left inferior parietal cortex (Rothi et al., 1991; Rushworth et al., 1997; Johnson-Frey, 2004). Furthermore, patients with apraxia of object use may be able to reach for and grasp objects normally (i.e., apraxia without optic ataxia).

The system that supports complex object-associated manipulations must be informed about the identity of the object, as object manipulations are specific to the identity and function of the object. This reasoning is in line with the pattern of findings that we have obtained: object representations in inferior parietal regions are contingent on analysis of the visual input by ventral structures. A key expectation of this view is that there will be privileged functional connectivity between inferior parietal HSF tool-preferring voxels and ventral stream structures that also exhibit tool preferences. We tested this expectation in a series of time-course based functional connectivity analyses. *ROI-based analysis of functional connectivity between the ventral and dorsal streams*

Several complementary analyses were carried out to test a key prediction made by the hypothesis that object representations in left inferior parietal regions are contingent on analysis of the visual input by ventral stream structures. In the first analysis, we localized the left medial fusiform gyrus within the ventral stream, which is known to be a tool-preferring region, with the data from Experiment 1 (broad-band stimuli; tools > animals; FDR q < .05). The resulting fusiform region of interest (ROI) is depicted in Figure 4A, and replicates previous observations using similar stimuli (e.g., Chao et al., 1999; Noppeney et al., 2006; Mahon et al., 2007). We then extracted the time series using data from Experiment 2 from *i*) the left medial fusiform ROI (Figure 4A), *ii*) the inferior HSF tool-preferring parietal region (Figure 2B), and the *ii*) the superior LSF tool-preferring parietal region (Figure 2B). Pearson correlation coefficients were used as measures of functional connectivity between each parietal region and the ventral stream ROI (after regression of noise confounds; see Materials and Methods above). There was greater functional connectivity between the left medial fusiform gyrus and the left inferior parietal ROI, than between the left medial fusiform and the left superior parietal ROI (t = 2.98, p < .02; paired, two-tailed; See Figure 4B). This analysis is conservative because a different group of subjects was used to define the ventral tool-preferring ROI.

We also repeated the analysis, defining the ventral tool-preferring ROI within Experiment 2, by collapsing across the factor spatial frequency and identifying voxels that responded more strongly to tool stimuli than animal stimuli (Contrast: Tools_(HSF + LSF) > Animals_(HSF + LSF); p < .005, corrected). The results of the connectivity analyses remained the same, indicating significantly greater connectivity between the inferior parietal HSF tool-preferring ROI and the ventral stream than between the superior parietal LSF tool-preferring ROI and the ventral stream (t = 2.49, p < .03).

In order to gain a deeper understanding of how connectivity to the two parietal ROIs was distributed across the population of voxels within the left medial fusiform gyrus, voxel-wise histograms of connectivity were plotted. As shown in Figure 4C, there is a shift in the distributions of connectivity strength within the left medial fusiform voxels: more voxels express stronger connectivity to the left inferior HSF tool-preferring

ROI than to the left superior LSF tool-preferring ROI. This difference was significant (p < .001), and the shift in the distributions was obtained regardless of whether the left fusiform was defined with the data from Experiment 1 (Figure 2B) or Experiment 2 (figure not shown; p < .001).

Whole brain analysis of functional connectivity.

In a second set of functional connectivity analyses, the superior and inferior parietal ROIs were used as seeds in whole-brain analyses of connectivity. At the same threshold (q < .05, FDR corrected) the left medial fusiform gyrus was identified by the analysis in which the inferior parietal HSF tool preferring ROI was used as a seed but not when the left LSF tool-preferring parietal ROI was used (Figure 5). This can be seen in the detailed depiction of the ventral stream (Figure 5A) in which voxels expressing connectivity to the left inferior parietal ROI are largely contained within the functionally defined left medial fusiform gyrus. Again, this correspondence was obtained both when the left medial fusiform ROI was defined with Experiment 1 and Experiment 2 (see Figure 5). In contrast, and constituting an important internal control, both parietal seeds identified the left middle temporal gyrus (lateral views, Figure 5).

Finally, we sought to test whether the spatial distribution of connectivity with inferior parietal regions matched the distribution of tool-preferences in the ventral stream. To that end, a lateral-to-medial analysis was carried out over a region of the ventral stream that straddled both medial and lateral areas. This analysis takes a swath of volume space, and 'slices' it into 3mm thick 'leaves', with each 'leaf' in a plane defined by Talairach z (superior-inferior) and Talairach y (anterior-posterior) coordinates (for precedent see, Mahon et al., 2009; see also Connolly et al., 2012). The statistic of

interest (e.g., contrast weighted t-values) are then averaged within the 'leaf' (i.e., along the anterior-posterior and inferior-superior dimensions), and plotted by their lateral-tomedial location (Talairach x-dimension). Two such lateral-to-medial analyses were carried out. The first was computed over the contrast weighted t-vales for the contrast of Tool_(HSE + LSE) > Animals_(HSE + LSE) in Experiment 2. Replicating previous lateral-to-medial analyses of contrast weighted statistics (Mahon et al., 2009; Connolly et al., 2012), tool preferences increase as the slices move more medial (blue line, Figure 6). The same lateral-to-medial analysis was then carried out over an index that represents the degree to which connectivity in the fusiform is differentially expressed to the inferior parietal ROI compared to the superior parietal ROI (IP = Inferior Parietal, SP = Superior Parietal: [connectivity with IP - connectivity with SP] / [connectivity with IP + connectivity SP]; see black line, Figure 6). The lateral-to-medial analysis of connectivity almost perfectly tracks the lateral-to-medial analysis of the contrast-weighted t-value. Specifically, as the position within the fusiform gyrus becomes more medial, tools lead to higher BOLD contrast than animals, and there is also differential connectivity to inferior parietal cortex compared to superior parietal regions. This high level of correspondence within the ventral stream between connectivity to parietal cortex and tool-preferences was observed regardless of whether the contrast-weighted t-map of tool preferences was derived from Experiment 1 (R^2 = .93, p < .01; Data not shown) or Experiment 2 (R^2 = .99, p < .001; as Shown in Figure 6).

Discussion

Models of conceptual representation in the human brain have been strongly influenced by the empirical fact that manipulable object stimuli, compared to a range of baselines, elicit differential BOLD contrast in regions of parietal cortex known to be involved in object prehension and use. Two theoretical inferences have been derived from that observation. The first is that the increased BOLD in parietal cortex reflects the automatic extraction of information about the way in which objects are typically manipulated by a dorsal visual route that is independent of the ventral stream. The second inference is that retrieval and simulation of the motor information involved in manipulating tools is a critical step in retrieving a tool concept from the visual presentation of a manipulable object. The findings that we have reported indicate that both assumptions need revision. The fact that tool preferences are observed in parietal cortex for stimuli that contain only HSF information indicates that at least some of the tool preferences in parietal cortex are contingent on analysis of the visual input by ventral stream structures. This conclusion is at odds with the hypothesis that recognition of manipulable objects is contingent on simulation of the motor movements associated with their use (for an overview, see Kiefer and Pulvermüller, 2012). This is because, if the motor simulation occurs only after the stimulus has been processed by ventral visual stream structures, then the stimulus has already gone through the classic channels of visual object identification. This conclusion converges with observations from brain damaged patients showing that damage to the left inferior parietal lobule can lead to an impairment for manipulating objects, while sparing the ability to recognize objects (for reviews, see Johnson-Frey, 2004; Mahon and Caramazza, 2008; Rothi et al., 1991). This conclusion also converges with prior arguments that the inferior parietal lobule and the functions it subserves should not be considered to be part of the dorsal visual pathway (Rizzolatti and Matelli, 2003; Milner and Goodale, 1995; Pisella et al., 2006; for

a parallel discussion of the right inferior parietal lobule, see Singh-Curry and Husain, 2009).

It was also observed that superior parietal regions show tool preferences for images that contain only LSF information. Neuropsychological data indicate that damage to superior parietal regions can lead to visuomotor impairments for reaching and grasping objects. Thus, within the context of the Embodied Cognition Hypothesis, it might be argued that the critical motor simulation is not of complex object manipulation (inferior parietal) but of the hand-shape configuration(s) that are implicated by object grasps. That direction for reconfiguring the proposal faces several difficulties. First, hand configurations appropriate for grasping objects will be shared by similarly shaped objects; thus, it is not at all clear that knowing the hand configuration appropriate for grasping an object could provide any individuating information that could contribute to distinguishing a stimulus from perceptual neighbors. Second, one of the original phenomena that motivated the proposal of a separation between ventral and dorsal visual streams was the dissociation between object prehension and object recognition (optic ataxia versus visual form agnosia). Thus, we know that patients with lesions to posterior-superior parietal cortex can be impaired for object prehension but spared for visual recognition (for discussion see Goodale and Milner, 1992; for a relevant conceptual analysis see Wu, 2008).

The hypothesis that the retrieval of complex object-associated manipulation knowledge is contingent on analysis of the visual input by the ventral visual pathway is not incompatible with the hypothesis that the retrieval of object use knowledge is automatic. However, if the activation of object use information is in fact automatic, and yet it is not necessary for recognition, then a cascading model of the dynamics of activation flow between concepts and object use information is strongly suggested (for discussion of dynamics as it relates to the embodied cognition hypothesis, see Mahon and Caramazza, 2008). Independent of these consideration about the dynamical principles about the concept-motor interface, our findings show that the automatic activation of parietal cortex when viewing tools does not monolithically represent processing of the visual input by the dorsal visual pathway. Differential BOLD responses to manipulable objects in parietal cortex reflect comingling of information extracted by the ventral stream with motor-based information represented in parietal regions, as well as the computation of reach and grasp information by the (classically understood) dorsal visual pathway.

One objection that may be raised is that the 'classic' construal of the function of the dorsal stream is itself insufficient. Recently the classic view of the division of labor between the dorsal and ventral visual streams has been questioned in light of evidence that object representations in posterior parietal regions exhibit response characteristics that generalize across image transformations (Konen and Kastner, 2008). Konen and Kastner's findings indicate that object representations in parietal cortex are abstracted away from the particular image characteristics of the stimuli, a property that may not be expected by the classic understanding of the dorsal stream. Those data, in the context of our current findings, outline a new framework for exploring whether there are mutual dependencies between the representations in the dorsal and ventral visual systems, and if so, over what types of information those dependencies are expressed. For instance, one open issue is whether regions of parietal cortex that show invariance in BOLD responses across image transformations derive their input entirely from a dorsal analysis of the visual input, or are also (in part) contingent on analysis of the visual input by ventral stream structures.

Another, and related, question that may be raised is whether the tool preferences for HSF stimuli in the left inferior parietal lobule are supported by the limited parvocellular channels that do putatively project to parietal cortex. In other words, perhaps parvocellular analysis of the visual input bypasses the ventral stream and subserves the tool preferences for HSF stimuli in the left inferior parietal lobule. One argument against this interpretation is the pattern of functional connectivity that we have reported between the left inferior parietal lobule and the left medial fusiform gyrus. Those functional connectivity increase the likelihood that the effects observed in the left inferior parietal lobule are in fact tied to processing in the ventral stream. Nevertheless, the broader point is that the functions of the classically understood dorsal visual pathway are functions largely attributable to magnocellular analysis (e.g., fast conductance, luminance based, sensitive to fast motion, etc.; e.g., Goodale and Milner, 1992; Merigan and Maunsell, 1993). Thus, even if the tool effects in the left inferior parietal lobule were driven by the small proportion of inputs to posterior parietal cortex that are parvocellular, then the inference would still be sanctioned that the representations in the left inferior parietal lobule are not part of the classically understood dorsal stream. This conclusion is consistent with the initial suggestions made by Goodale and Milner (1992).

The stronger conclusion, and which we favor is that the tool preferences in the left inferior parietal lobule are in fact contingent on analysis of the visual input by the ventral object processing stream. A critical test of this stronger conclusion would be provided by studying the distribution of tool preferences in parietal cortex in patients with lesions to the ventral stream. The framework that we have outlined predicts that patients with ventral stream lesions affecting the visual and semantic analysis of manipulable object stimuli will continue to show normal tool responses in posterior and superior parietal regions, but will have attenuated or even abolished responses in the left inferior parietal lobule.

In summary, and together with prior neuropsychological research, our findings suggest an alternative to the Embodied Cognition Hypothesis of tool recognition: the activation of (at least some) motor-relevant information in parietal cortex is contingent on processing within the ventral stream. On this view, motor-relevant information about object use is not a necessary intermediary step in object recognition (Machery, 2007; Mahon and Caramazza, 2008; Chatterjee, 2010). An interesting issue for future research is whether previously described white matter connectivity between the temporal and parietal lobules (Zhong and Rockland, 2003; Rushworth et al., 2006) is the basis for the integration of information about object identity with information about object-associated manipulations.

Authors' Note

This research was supported by NIH grant R21 NS076176 from NINDS. JA was supported by Fundação para a Ciência e a Tecnologia fellowship SFRH/BPD/70970/2010 and by EU-FP7/2007-2013, under PCOFUND-GA-2009-246542. This research was supported in part by Norman and Arlene Leenhouts. The authors are grateful to Jessica Cantlon, Greg DeAngelis, and Bill Merigan for discussion.

Figure Captions

Figure 1. Examples of stimuli used in Experiments 1 and 2. The same set of grey scale tools and animals were used across both experiments. The images were broad-band in Experiment 1, and filtered to contain only high or low spatial frequencies in Experiment 2.

Figure 2. Tool preferring regions of left parietal cortex dissociate according to the spatial frequency content of the images. **Panel A.** Detail of left parietal tool-preferring voxels in Experiment 1 (FDR q < .05) obtained using standard (broad-band) grey scale images. **Panel B**. Detail of the left parietal regions defined by the contrasts of Tools_{HSF} > Animals_{HSF} (purple) and Tools_{LSF} > Animals_{LSF} (green) (both maps p < .05, corrected). These contrasts, which are independent and so have no bias against showing overlap, indicate a dissociation along the inferior-to-superior dimension in whether the tool-preference is differentially carried by HSF or LSF information in the images.

Figure 3. Overlap of contrast maps for Experiments 1 and 2 in left parietal cortex. Plotted on the orange color scale is the contrast of Tool > Animal (FDR, q < .05) from Experiment 1. Plotted on the purple color scale is the contrast Tool_{HSF} > Animal_{HSF}, and plotted on the green color scale is the contrast Tool_{LSF} > Animal_{LSF}, both from Experiment 2 (p < .05, corrected). Within left parietal cortex, more inferior aspects show tool preferences for high spatial frequency images, while more superior regions exhibit tool preferences for low spatial frequency images. **Figure 4**. The inferior HSF tool-preferring parietal region exhibits privileged connectivity with the left medial fusiform gyrus. **Panel A.** The medial fusiform ROI from Experiment 1 is shown, defined by the contrast Tools > Animals (FDR q < .05). **Panel B.** Time-course based functional connectivity analyses were conducted using time-series data from the left medial fusiform ROI (Panel A), and the inferior and superior parietal ROIs (Figure 2B). There was significantly greater connectivity between the left medial fusiform ROI and the inferior parietal ROI compared to the superior parietal ROI. Error bars reflect standard error of the means across subjects. **Panel C.** Voxel-wise connectivity values (t-values for the group-level analysis) of connectivity with the two parietal ROIs were extracted for each voxel within the left medial fusiform gyrus, and plotted as histograms. The results indicate a shift in the distributions, with stronger connectivity between the medial fusiform voxels and the inferior parietal ROI, compared to the superior parietal ROI.

Figure 5. Whole-brain functional connectivity maps computed using the inferior and superior parietal ROIs (from Figure 2). Group-level statistics were computed as one-sample ttests (voxel-wise) over subjects' connectivity values. **Panel A.** Group-level t-map for connectivity with the left inferior HSF tool-preferring parietal ROI. The detail shows that the connectivity analysis identifies voxels in the left medial fusiform gyrus, as functionally defined with either Experiment 1 (red border) or Experiment 2 (blue border). **Panel B.** Group-level t-map for connectivity with the left superior LSF tool-preferring parietal ROI. As shown in the detailed image, connectivity with the superior parietal ROI.

is absent within the independently defined medial fusiform ROIs. All t-maps were thresholded at FDR q < .05.

Figure 6. Lateral-to-medial analysis of ventral stream tool-preferences and connectivity with parietal cortex. Two independent lateral-to-medial analyses were computed by averaging the relevant statistic along the Talairach-z (inferior-superior) and Talairach-y (anterior-posterior) axes for each Talairach-x dimension value (binned at 3mm). In one analysis, the lateral-to-medial index for the contrast weighted t-value measuring tool preferences (Tools > Animals) was computed (light blue line). In a second lateral-to-medial analysis, differential connectivity to the left inferior parietal ROI was determined: (Connectivity with inferior parietal ROI – Connectivity with superior parietal ROI) / (Connectivity with inferior parietal ROI + Connectivity with superior parietal ROI). The results of the two independent analyses demonstrate correspondence along the lateral-to-medial dimension.

References

- Barsalou LW (1999) Perceptual symbol systems. Behav Brain Sci 22:577-609; discussion 610-560.
- Brainard DH (1997) The Psychophysics Toolbox. Spat Vis 10:433-436.
- Buccino G, Riggio L, Melli G, Binkofski F, Gallese V, Rizzolatti G. 2005. Listening to action related sentences modulates the activity of the motor system: a combined TMS and behavioral study. *Cogn. Brain Res.*, 24, 355–63
- Chao LL, Martin A (2000) Representation of manipulable man-made objects in the dorsal stream. NeuroImage 12:478-484.
- Chao LL, Haxby JV, Martin A (1999) Attribute-based neural substrates in temporal cortex for perceiving and knowing about objects. Nat Neurosci 2:913-919.
- Connolly, A.C., Guntupalli, J.S., Gors, J., Hanke, M., Halchenko, Y.O., Wu, Y-C., Abdi,
 H., Haxby, J.V. (2012). The representation of biological classes in the human brain. *The Journal of Neuroscience*, 32, 2608-2618.

Chatterjee A (2010) Disembodying Cognition. Language and Cognition 2:79-116.

- Derrington AM, Lennie P, Journal of Physiology (London) (1984) Spatial and temporal contrast sensitivities of neurones in lateral geniculate nucleus of macaque. 357:219-240.
- Felleman DJ, Van Essen DC (1991) Distributed hierarchical processing in primate visual cortex. Cerebral Cortex, 1:1-47.
- Ferrera VP, Nealey TA, Maunsell JHR (1992) Mixed parvocellular and magnocellular geniculate signals in visual area V4. . Nature 358:756-758.

- Gallese V, Lakoff G (2005) The brain's concepts: The role of the sensory-motor system in conceptual knowledge. Cognitive Neuropsychology 22:455–479.
- Goodale MA, Milner AD (1992) Separate visual pathways for perception and action. *Trends in Neurosciences* 15:20-25.
- Johnson-Frey SH (2004) The neural bases of complex tool use in humans. Trends in Cognitive Sciences 8:71-78.
- Kiefer, M. and Pulvermüller, F. (2012). Conceptual representations in mind and brain: Theoretical developments, current evidence and future directions. *Cortex*, 48, 805-825.
- Konen CS, Kastner S (2008) Two hierarchically organized neural systems for object information in human visual cortex. Nature neuroscience 11:224-231.
- Livingstone M, Hubel D (1988) Segregation of form, color, movement, and depth: anatomy, physiology, and perception. Science 240:740-749.

Machery E (2007) Concept empiricism: A methodological critique. Cognition 104:19-46.

- Mahon BZ, Caramazza A (2008) A critical look at the embodied cognition hypothesis and a new proposal for grounding conceptual content. J Physiol Paris 102:59-70.
- Mahon BZ, Milleville SC, Negri GA, Rumiati RI, Caramazza A, Martin A (2007) Actionrelated properties shape object representations in the ventral stream. Neuron 55:507-520.
- Mahon BZ, Anzellotti S, Schwarzbach J, Zampini M, Caramazza, A. (2009). Categoryspecific organization in the human brain does not require visual experience. *Neuron*, 63, 397-405.

- Martin A, Ungerleider LG, Haxby JV. 2000. Category specificity and the brain: the sensory/motor model of semantic representations of objects. In Higher Cognitive Functions: The New Cognitive Neurosciences, ed. MS Gazzaniga, pp. 1023–36. Cambridge, MA: MIT Press
- Maunsell JHR, Nealey TA, DePriest DD (1990) Magnocellular and parvocellular contributions to responses in the middle temporal visual area (MT) of the macaque monkey. Journal of Neuroscience 10:3323-3334.
- Merigan WH, Maunsell JH (1993) How parallel are the primate visual pathways? Annu Rev Neurosci 16:369-402.
- Milner A.D., Goodale M.A. 1995. The visual brain in action. Oxford University Press.
- Noppeney U, Price CJ, Penny WD, Friston KJ (2006) Two distinct neural mechanisms for category-selective responses. Cereb Cortex 16:437-445.
- Pelli DG (1997) The VideoToolbox software for visual psychophysics: Transforming numbers into movies. Spatial Vision 10:437-442.
- Pisella, L., Binkofski, F., Lasek, K., Toni, I., and Rossetti, Y. (2006). No doubledissociation between optic ataxia and visual agnosia: Multiple sub-streams for multiple visuo-manual integrations. *Neuropsychologia*, 44, 2734-2748.
- Pulvermuller F, Hauk O, Nikulin VV, Ilmoniemi RJ (2005) Functional links between motor and language systems. European Journal of Neuroscience 21:793-797.
- Rizzolatti, G. and Matelli, M. (2003). Two different streams form the dorsal visual system: anatomy and functions. *Experimental Brain Research*, 153, 146-157.
- Rothi LJG, Ochipa C, Heilman KM (1991) A Cognitive Neuropsychological Model of Limb Praxis. Cognitive Neuropsychology 8:443-458.

- Rushworth MF, Behrens TE, Johansen-Berg H (2006) Connection patterns distinguish 3 regions of human parietal cortex. Cerebral cortex 16:1418-1430.
- Rushworth MF, Nixon PD, Renowden S, Wade DT, Passingham RE (1997) The left parietal cortex and motor attention. Neuropsychologia 35:1261-1273.
- Schwarzbach J (2011) A simple framework (ASF) for behavioral and neuroimaging experiments based on the psychophysics toolbox for MATLAB. Behavior research methods.
- Singh-Curry, V. and Husain, M. (2009). The functional role of the inferior parietal lobe in the dorsal and ventral stream dichotomy. *Neuropsychologia*, 47, 1434-1448.
- Talairach J, Tournoux P (1988) Co-Planar Stereotaxic Atlas of the Human Brain. New York: Thieme.
- Tootell RB, Silverman MS, Hamilton SL, Switkes E, De Valois RL (1988) Functional anatomy of macaque striate cortex. V. Spatial frequency. J Neurosci 8:1610-1624.
- Ungerleider LG, Mishkin M (1982) Two cortical visual systems. In: Analysis of Visual Behavior (Goodale MA, Mansfield RJW, eds), pp 549-586. Cambridge: The MIT Press.
- Wu WC (2008) Visual attention, conceptual content, and doing it right. Mind 117:1003-1033
- Zhong Y-M, Rockland KS (2003) Inferior parietal lobule projections to anterior inferiortemporal cortex (area TE) in Macaque monkey. Cerebral Cortex 13:527-

Figure 1. Examples of Stimuli Used in Experiments 1 and 2



Figure 2. Statistical Contrast Maps for Tool Preferences in Parietal Cortex

A. Experiment 1. Tool-preferences for broad band stimuli



B. Experiment 2. Tool-preferences dissociate for HSF and LSF images



Figure 3. Contrast Maps from Experiments 1 and 2 Overlaid



Figure 4. Parietal-Fusiform Connectivity

A. Left Medial Fusiform ROI

B. Connectivity between fusiform and parietal ROIs



C. Distributions of connectivity with parietal ROIs for voxels in the Left Medial Fusiform ROI



Figure 5. Whole Brain Connectivity with Left Parietal Tool Preferring ROIs A. The HSF tool-preferring parietal seed identifies the left medial fusiform



B. The LSF tool-preferring parietal seed does not identify the left medial fusiform



Figure 6. Lateral-to-medial correspondence in the ventral stream between tool-preferences and connectivity to parietal ROIs

