

DRD2 polymorphism and sensitivity to losses during value-based decision-making


Cristina Bañuelos^{1,3}, Kasey Creswell, Ph.D.^{1,3}, Stephen Manuck, Ph.D.^{2,3}, Peter Gianaros, Ph.D.^{2,3}, Timothy Verstynen, Ph.D.^{1,3,4}

1. Department of Psychology, Carnegie Mellon University
2. Department of Psychology, University of Pittsburgh School of Medicine
3. Neuroscience Institute, Carnegie Mellon University and University of Pittsburgh
4. Department of Biomedical Engineering, Carnegie Mellon University



Background


- Dopamine receptor subtypes respond differently to gains and losses
- Dopamine receptor D2 (DRD2) -141C Ins/Del genetic variant associates with lower levels of D2 receptor density
- DRD2 variant carriers may be less sensitive to losses during feedback learning
- It remains unclear how phasic dopamine is modulated by the opposing influences on the direct and indirect striatal pathways in feedback learning



Hypothesis

- Ventral striatal reactivity to rewards interacts with the presence of the DRD2 -141C Ins/Del variant to impact sensitivity to gains and losses


Results: Distribution of Measures



- Distributions of Age, VS reactivity and IGT scores by DRD2 group are not normal

Results: VS Reactivity between Groups

- DRD2 variant carriers had generally lower performance in IGT than non-carriers for Payoff ($t = 3.23$, $p = 0.001$), but not for Sensitivity ($t = 0.54$, $p = 0.590$), and lower VS reactivity ($t = -1.77$, $p = 0.077$)



Results: Main Effects and Interactions

Payoff Analysis

- Significant effects from the DRD2 variant and for VS reactivity, but not for the interaction
- Carriers scored almost 9 points lower on Payoff compared to the non-carriers
- Strong evidence against the alternate model

Payoff Models

Bayes Factor: 18.37

$$\begin{aligned} H_0: Y_P &= \hat{\beta}_0 + \hat{\beta}_1 X_{DRD2} + \hat{\beta}_2 X_{VS} \\ H_A: Y_P &= \hat{\beta}_0 + \hat{\beta}_1 X_{DRD2} + \hat{\beta}_2 X_{VS} + \hat{\beta}_3 X_{DRD2} \times X_{VS} \end{aligned}$$

Payoff Linear Regression:

	coef	std err	t	P> t	[0.025	0.975]
Intercept	16.7039	2.020	8.269	0.000	12.733	20.674
DRD2	-8.7905	3.634	-2.419	0.016	-15.932	-1.649
VS	21.1400	10.020	2.110	0.035	1.447	40.833
DRD2:VS	-9.5730	18.830	-0.508	0.611	-46.582	27.436

Sensitivity Linear Regression:

	coef	std err	t	P> t	[0.025	0.975]
Intercept	30.6606	1.746	17.563	0.000	27.230	34.092
DRD2	1.7348	3.140	0.553	0.581	-4.437	7.906
VS	4.7830	8.658	0.552	0.581	-12.234	21.801
DRD2:VS	-1.3082	16.272	-0.080	0.936	-33.289	30.673

Sensitivity Analysis

- No significant effects from the DRD2 variant, for VS reactivity, nor for the interaction
- Strong evidence against the alternate model

Sensitivity Models


Bayes Factor: 20.86

$$\begin{aligned} H_0: Y_Q &= \hat{\beta}_0 + \hat{\beta}_1 X_{DRD2} + \hat{\beta}_2 X_{VS} \\ H_A: Y_Q &= \hat{\beta}_0 + \hat{\beta}_1 X_{DRD2} + \hat{\beta}_2 X_{VS} + \hat{\beta}_3 X_{DRD2} \times X_{VS} \end{aligned}$$

Methods

Participants

- Neurologically healthy adults from southwestern Pennsylvania from University of Pittsburgh's Adult Health And Behavior project, Phase II
- 438 participants (228 females, 210 males) between the ages of 30 and 54 ($M=42.67$, $SD=7.36$)
- Participant's blood was drawn for genotyping for the DRD2 -141C Ins/Del variant (119 carriers, 319 non-carriers) (1)



Iowa Gambling Task (IGT)

- Maximize profit with a set amount of number of draws possible from 4 decks of cards

Payoff:

$$P = (C + D) - (A + B)$$

Frequency of Reward:


Deck A	Deck B	Deck C	Deck D
Low	High	Low	High
Loss	Loss	Gain	Gain

Sensitivity to high frequency rewards:

$$Q = (B + D) - (A + C)$$

Ventral Striatal (VS) Reactivity Task

- Assess the VS reactivity to negative and positive feedback cues associated with monetary gain (2 & 3)



Conclusion

Summary

- Lower D2 receptor density, and lower dopamine release associates with sub-optimal value-based decision-making
- The interaction between D2 receptor density and dopamine release does not associate with differences in value-based decision-making
- No significant effects were found for sensitivity to high frequency rewards

Take Home Message

- Strong evidence that lower available D2 receptors, and lower dopamine release associates with a lower sensitivity to gains and losses in feedback learning
- Inconclusive evidence for the mechanistic role of the D2 pathway in using feedback effectively during value-based decision-making

Future Directions

Theoretical Directions

- Simulations of the D1 and D2 pathways in order to test other possible mechanistic roles of the D2 pathway in feedback learning

Experimental Directions

- Another feedback learning measure that separates between sensitivity to losses and gains

References

- Lerman, C., Jepson, C., Wileyto, E. P., Epstein, L. H., Rukstalis, M., Patterson, F., ... & Berrettini, W. (2006). Role of functional genetic variation in the dopamine D2 receptor (DRD2) in response to bupropion and nicotine replacement therapy for tobacco dependence: results of two randomized clinical trials. *Neuropsychopharmacology*, 31(1), 231-242.
- Hariri, A. R., Brown, S. M., Williamson, D. E., Flory, J. D., De Wit, H., & Manuck, S. B. (2006). Preference for immediate over delayed rewards is associated with magnitude of ventral striatal activity. *Journal of Neuroscience*, 26(51), 13213-13217.
- Gianaros, P. J., Manuck, S. B., Sheu, L. K., Kuan, D. C., Votruba-Drzal, E., Craig, A. E., & Hariri, A. R. (2011). Parental education predicts corticostriatal functionality in adulthood. *Cerebral Cortex*, 21(4), 896-910.